

PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification 6: A61K 31/70, A61F 13/20, G02C 7/02	A1	(11) International Publication Number: WO 95/07085 (43) International Publication Date: 16 March 1995 (16.03.95)
(21) International Application Number: PCT/US94/10175 (22) International Filing Date: 7 September 1994 (07.09.94) (30) Priority Data: 08/116,908 7 September 1993 (07.09.93) US (71) Applicant: ESCALON OPHTHALMICS, INC. [US/US]; 182 Tamarack Circle, Skillman, NJ 06558 (US). (72) Inventor: BENEDETTO, Dominick, A.; 124 Avenue B, Bayonne, NJ 07002 (US). (74) Agent: SAUNDERS, Thomas, M.; Lorusso & Loud, 440 Commercial Street, Boston, MA 02109 (US).		(51) Designated States: CA, JP, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE). Published <i>With international search report.</i>

(54) Title: SURFACE ACTIVE VISCOELASTIC SOLUTIONS FOR OCULAR USE**(57) Abstract**

This invention encompasses a modified mucopolysaccharide solution for use as a biologically active therapeutic infusion comprising a pharmaceutical grade viscoelastic fraction selected from a group consisting of an acyl-substituted hyaluronic acid having acyl groups thereof with three to twenty carbon atoms and mixtures of said acyl-substituted hyaluronic acid with hyaluronic acid, and hydroxypropylmethylcellulose. In particular these solutions have a surface tension of between 40 and 65 dynes/cm²; particularly a viscoelastic fraction has an average molecular weight of at least 50,000. In some embodiments a physiological buffer fraction is present. This invention further encompasses a method of using the claimed composition.

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	GB	United Kingdom	MR	Mauritania
AU	Australia	GE	Georgia	MT	Malta
BB	Barbados	GN	Guinea	NE	Niger
BE	Belgium	GR	Greece	NL	Netherlands
BF	Burkina Faso	HU	Hungary	NO	Norway
BG	Bulgaria	IE	Ireland	NZ	New Zealand
BJ	Benin	IT	Italy	PL	Poland
BR	Brazil	JP	Japan	PT	Portugal
BY	Belarus	KE	Kenya	RO	Romania
CA	Canada	KG	Kyrgyzstan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SI	Slovenia
CI	Côte d'Ivoire	LI	Liechtenstein	SK	Slovakia
CM	Cameroon	LK	Sri Lanka	SN	Senegal
CN	China	LU	Luxembourg	TD	Chad
CS	Czechoslovakia	LV	Latvia	TG	Togo
CZ	Czech Republic	MC	Monaco	TJ	Tajikistan
DE	Germany	MD	Republic of Moldova	TT	Trinidad and Tobago
DK	Denmark	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	US	United States of America
FI	Finland	MN	Mongolia	UZ	Uzbekistan
FR	France			VN	Viet Nam
GA	Gabon				

WO 93/07085

PCT/US94/10175

1 SURFACE ACTIVE VISCOELASTIC SOLUTIONS FOR OCULAR USE

2

3 This application is a continuation-in-part of copending
4 U.S. Pat. App. 08/061,773 filed May 13, 1993, which is a
5 continuation of U.S. Pat. App. 07/440,078 filed November 22,
6 1989, now abandoned.

7

8 Field of the Invention.

9 The present invention relates to ophthalmic solutions for
10 use during ocular and intraocular surgery, and more particularly
11 to the use of surface active viscoelastic solutions during the
12 extraction of a cataractous human lens and the implantation of a
13 prosthetic ocular and intraocular lens. During surgery, the use
14 of ophthalmic infusions with controlled physical properties,
15 especially surface activity and viscoelastic properties, is
16 advantageous for (1) replacing the fluid aqueous humor or ocular
17 and intraocular air, (2) protecting the internal structures of
18 the eye from accidental instrument or ocular and intraocular
19 prosthetic device contact, (3) preventing irrigation damage by
20 solutions used in routine cataract surgery, and (4) retarding
21 aspiration from the eye of the viscoelastic solution during the
22 surgical procedure. In addition, the invention relates to a
23 method of adhering a contact lens to the surface of the eye,
24 such as in association with procedures permitting a medical
25 professional to view ocular and intraocular structures through
26 the contact lens and through the viscoelastic solution. In

27

28

WO 95/07085

PCT/US94/10175

1 another application, the viscoelastic solution of this invention
2 is used by injecting the solution into or under tissues within
3 the eye, such as to dissect tissue off of the retina.

4 Background of the Invention

5 In the past, biocompatible polymers used in ocular and
6 intraocular surgery have been the naturally occurring
7 mucopolysaccharides hyaluronic acid and chondroitin sulfate;
8 mixtures of hyaluronic acid and chondroitin sulfate; and,
9 cellulose derivatives, such as hydroxypropylmethylcellulose
10 (HPMC). Table 1
11 presents data reported in Viscoelastic Materiale, Ed. E.S.
12 Rosen, Proceedings of the Second International Symposium of the
13 Northern Eye Institute, Manchester [U.K.], 17-19 July, 1986
14 (Pergamon Press, New York) as to the molecular weight of
15 commercially available ocular products. Depending on the source
16 from which these mucopolysaccharides are drawn, the molecular
17 weights are estimated in the 50,000 range with the hyaluronic
18 acid extending upwards to the 8×10^6 range. Hyaluronic acid
19 was first isolated and characterized by Meyer, Palmer and
20 reported in the J. Biol. Chem., Vol. 107, p. 629 (1934) and Vol.
21 114, p.689 (1936) and by Balazs in the Fed. Proc. Vol. 17, p.
22 1086 (1958); and chondroitin sulfate by Bray et al. in Biochem.
23 J. Vol. 38, p. 144 (1944); and Patat, Elias, Z. Physiol. Chem.
24 vol. 316, p. 1 (1959).

25
26 Literature in the art describes the basic isolation and
27 characterization of the viscoelastic solutions. It is a
28 surprising feature of this invention which describes the control

WO 95/07085

PCT/US94/10175

1 of viscoelastic properties as related to the surface activity,
2 or the solution fracturing under applied stress. In particular,
3 it is surprising to manipulate or enhance the physical
4 properties of viscoelastic solutions of mucopolysaccharides,
5 hyaluronic acid, and/or chondroitin sulfate. It is believed
6 that disclosure here of a processes to provide hyaluronic acid
7 and species thereof with controlled surface activity is unique.
8 This is also especially true of the control of surface activity
9 of mucopolysaccharide solutions by the addition of biologically
10 compatible surfactants. A characteristic feature of
11 biologically compatible surfactants is the absence of observed
12 alteration in cellular physiology upon contact. Early work in
13 the viscoelastic field was presented by the inventor of this
14 disclosure and his associates. Benedetto, D.A. et. al.,
15 Viscoelastic Materials: Basic Science and Clinical Application.
16 (Symposium Proceedings), University of Manchester, England, July
17 17-19, 1986.

18
19 As to commercial production, a review of the ophthalmic
20 pharmacopoeia reveals there are several viscoelastic solutions
21 produced for ocular and intraocular use during ophthalmic
22 surgery. The most common application for these solutions is in
23 the intraocular lens implant procedure for human cataract
24 surgery. This procedure involves extraction of the cataractous
25 human lens through a small surgical opening in the eye and the
26 replacement of the lens by a prosthetic intraocular lens placed
27 in situ. Biocompatible polymers presently or previously in use
28 are hyaluronic acid (Healon™, Amvisc™); chondroitin sulfate, and

WO 95/07085

PCT/US94/10175

1 a combined solution of hyaluronic acid and chondroitin sulfate
2 (Viscoat™); and a hydroxypropylmethylcellulose solution
3 (Occucoat™). Research conducted recently demonstrates that
4 Healon™ and Amvisc™ are not surface active, but Viscoat™ and
5 Occucoat™ are.

6 Chondroitin sulfate does not exist as a free polysaccharide
7 in its native state, but as a proteoglycan. It is obtained from
8 sources associated with protein contaminants. The avoidance of
9 chondroitin sulfate avoids a potential source of pyrogenic
10 reaction, and the substantial cost associated with protein
11 removal.

12 Summary of the Invention

13 The invention presented herein discloses modified
14 mucopolysaccharide or viscoelastic solutions for use as
15 biologically active therapeutic infusions. In one form of the
16 invention, the mucopolysaccharide solution is formed from a
17 viscoelastic fraction and a buffer fraction. It has been found
18 that when a new synthetic molecule acyl-substituted hyaluronic
19 acid is employed as the viscoelastic fraction, control of
20 surface activity is achieved. An indicia of this is the
21 decrease of the surface tension of the solution which is now
22 within predetermined limits discussed below. Surface tension
23 modification is also accomplished with viscoelastic fractions in
24 which the acyl-substituted hyaluronic acid is mixed with one or
25 more of hyaluronic acid; and hydroxypropylmethylcellulose. In
26 certain applications, the viscoelastic solution of this
27 invention is used in a method of adhering a contact lens to the
28

WO 95/07085

PCT/US94/10175

1 surface of the eye, such as in association with procedures
2 permitting a medical professional to view ocular and intraocular
3 structures through the contact lens and through the viscoelastic
4 solution. This is particularly useful in facilitating surgical
5 procedures. In another application, the viscoelastic solution of
6 this invention is used by injection the solution into or under
7 structures or tissues within the eye, such as to dissect tissue
8 off of the retina.

9
10 In the broadest terms, surface active viscoelastic
11 solutions with controlled solution properties, are characterized
12 by surface tension, equilibrium contact angle, dynamic
13 viscosity, and cohesiveness (the measure of solution fracture
14 under stress). In a particular embodiment, this invention is
15 delimited by the three dimensional representation of Fig. 7.

16 In one example, bioengineered hyaluronic acid from a
17 bacterial source with an average molecular weight of 50,000 is
18 modified by acyl substitution with three to twenty carbon atom
19 acyl groups so that the resultant surface tension of such a
20 solution is between 40 and 65 dynes/cm². In the practice of
21 this invention, a viscoelastic solution having a surface tension
22 of less than about 56 dynes/cm², and more particularly, less
23 than about 50 dynes/cm² is of particular advantage.

24
25 This invention comprises a modified mucopolysaccharide
26 solution for use as a biologically active therapeutic infusion
27 comprising:

28

WO 95/07085

PCT/US94/10175

BEST AVAILABLE COPY

1 a pharmaceutical grade viscoelastic fraction selected from
2 the group consisting of acyl-substituted hyaluronic acid having
3 acyl groups thereof with three to twenty carbon atoms,
4 hyaluronic acid, hydroxypropylmethylcellulose and mixtures
5 thereof, and absent chondroitin sulfate said fraction having a
6 surface tension of between 40 and 65 dynes/cm²; and,

7 optionally with a physiological buffer fraction, such that
8 the viscoelastic comprises about a 0.1% percent of the solution
9 to about 5% of the solution, by weight, and preferably from
10 about 0.5 % to about 3%;

11 said modified mucopolysaccharide solution having a
12 viscosity of between 10,000 and 100,000 centipoise when measured
13 at a shear rate of 3 sec⁻¹ at 25°C; and,

14 optionally wherein the modified mucopolysaccharide
15 solution has a surface tension of less than about 56 dynes/cm²,
16 and further a surface tension of less than about 50 dynes/cm²;
17 and further,

18 optionally wherein the solution has an osmolality of from
19 about 250 to about 400 milliosmoles, or is generally isotonic
20 with ophthalmic tissue.

21 In some embodiments the modified mucopolysaccharide
22 solution viscoelastic fraction has an average molecular weight
23 of at least 50,000. Reference is further made to the
24 viscoelastic fraction being an acyl-substitute hyaluronic acid
25 having acyl groups thereof with three to twenty carbon atoms.

26 In particular applications the modified mucopolysaccharide
27 solution of this invention includes a surfactant fraction of a
28 biocompatible component selected from a group consisting of